

OUR FORMULA IS:  
HELP YOUR FARM  
GROW HEALTHY

Constant investments in technology to create a wide line of high-quality products, helping the poultry farmer to obtain greater productivity and profitability.

Provide the animals on your farm with the correct dose of medication, using the Calculation of Medication Via Water, which can be found on our website:  
[www.sauvet.com.br/calculo\\_agua.php](http://www.sauvet.com.br/calculo_agua.php)



ATTENTION IN THE PRE AND  
POST SALE.



DEVELOPMENT OF  
INNOVATIVE PRODUCTS.



AVAILABILITY AND  
IMMEDIATE DELIVERY.



COMMITMENT TO THE  
PERFORMANCE OF YOUR FARM



FOOD SAFETY TO THE FINAL  
CONSUMER



100% BRAZILIAN COMPANY



**MAXIMULIN PS**



#### PACKING

Sachets of 250 g in boxes of 5 Kg.



#### COMPOSITION

Each 100g contains:

Tiamulin (Hydrogen fumarate)..... 45g

Excipient (enough. For) .....100g



#### ADMINISTRATION

For oral administration: mix with the feed or diluted in drinking water.



#### INDICATION

MAXIMULIN OS it is indicated for the control of respiratory infections such as: Pleuropneumonia (*Actionobacillus leuropneumoniae*), Enzootic pneumonia (*Mycoplasma hyopneumoniae*), Pneumonia (*Pasteurella sp.*, *Haemophilus parasuis* and *Streptococcus spp.*); Gastrointestinal infections; Dysentery (*Corynebacterium sp.*, *Serpulina hyodysenteriae*), Ileitis (*Lawsonia intracellularis*), Colitis (*Serpulina pilosicoli*); Arthritis (*Mycoplasma hyosynoviae* and *Mycoplasma hyorhinis*).



#### DOSAGE

<i>It is indicated Dose per kg</i>	<i>Dose per kg live weight</i>
Treatment of blood dysentery	8mg
Treatment of ileitis and spirochetal colitis	10mg
Treatment of enzootic pneumonia	15mg



#### WAITING PERIOD

Suspend the use of the medication 10 days before the slaughter of animals intended for human consumption.



## MAXIMULIN 80



### PACKING

Sachet of 200 g in boxes of 5 kg and multi-layer bags containing 5kg and 25kg.



### COMPOSITION

Each 100g contains:  
Tiamulin (Hydrogen Fumarate).....80g  
Excipient. (enough for) .....100g



### ADMINISTRATION

For oral administration mix with the ration and dilute in drinking water.



### INDICATION

MAXIMULIN 80 is indicated for the treatment of respiratory and enteric diseases in pigs caused by *Streptococcus suis*.



### DOSAGE

Administer 5mg to 10mg of Tiamulin fumarate per Kg of live weight, for 10 consecutive days.



### WAITING PERIOD

Suspend the use of the medication 2 days before the slaughter of animals intended for human consumption.



## MAXIMULIN 10% PREMIX



### PACKING

Sacos multifolheados contendo 25Kg.



### COMPOSITION

Each 100g contains:

Tiamulin (Hydrogen Fumarate).....10g  
Excipient. (enough for) .....100g



### ADMINISTRATION

For oral administration: mix with the feed.



### INDICATION

MAXIMULIN 10% PREMIX IT is indicated for the treatment of respiratory diseases caused by *Mycoplasma spp.* and *Haemophilus spp.* and enteric diseases caused by *Serpulina hyodysenteriae* and *Campylobacter app.*



### DOSAGE

Provide 100 to 200ppm of Tiamulin per ton of feed for 5 to 10 consecutive days.



### WAITING PERIOD

Suspend the use of the medication 7 days before the slaughter of animals intended for human consumption.





## PACKING

Sachet of 1Kg in boxes of 10Kg and multilayer bags containing 25Kg.



## COMPOSITION

Each 100g contains:

Oxytetracycline Hydrochloride .....10g  
Sulfadiazine ..... 7,5g  
Trimethoprim .....1,5g  
Excipient (enough. For) .....100g



## ADMINISTRATION

For oral administration: mix with the feed.



## INDICATION

MEDMAXI is indicated for the treatment of infections caused by germs sensitive to the components of the formula, such as: Pneumonias (*Pasteurella sp*, *Bordetella sp*, *Haemophilus sp*, *Mycoplasma hyopneumoniae*), Pleuropneumonia (*Actinobacillus pleuropneumoniae*); Atrophic rhinitis (*Pasteurella multocida*, *Bordetella bronchiseptica*); Pos-weaning diarrhea (*Salmonella sp*, *Clostridium sp*, *Escherichia coli*); Cervical and joint abscesses (*Staphylococcus sp*, *Corynebacterium sp*, *Streptococcus sp*.)



## DOSAGE

Administer 3kg to 4kg of MEDMAXI per ton of feed for 5 to 7 consecutive days. It is advisable to continue treatment for 48 hours after the disappearance of symptoms.



## WAITING PERIOD

Suspend the use of the medication 18 days before the slaughter of animals intended for human consumption.



**MAXNOR 50**



#### PACKING

Sachets of 100g in boxes of 5kg and multilayer bags containing 25kg.



#### COMPOSITION

Each 100g contains:  
Norfloxacin base..... 50g  
Excipient (enough. For) .....100g



#### ADMINISTRATION

For oral administration: mix with the feed.



#### INDICATION

MAXIMOR 50 it is indicated for the control of bacterial infections, such as: Enzootic pneumonia: *Mycoplasma hyopneumoniae*; Pneumonia: *Pasteurella multocida*, *Mannheimia haemolytica*, *Bordetella bronchiseptica*, *Haemophilus pleuropneumoniae*. Colibacillosis: *E. coli*; MMA syndrome (metritis-mastitis-agalaxia): *Staphylococcus aureus*, *Streptococcus agalaciae*, *Streptococcus dysgalactiae*; *Staphylococci*: *Staphylococcus aureus*; Streptococci: *Streptococcus ssp*.



#### DOSAGE

Give 0.5mg to 10mg of norfloxacin per kg of body weight for 3 to 5 consecutive days. For salmonellosis, administer for 5 consecutive days. Continue with the treatment for 48 hours after the disappearance of the symptoms or at the discretion of the veterinarian.



#### WAITING PERIOD

Suspend the use of the medication 4 days before the slaughter of animals intended for human consumption.





  
**CIPROMAX**



#### PACKING

Sachets of 100g in boxes of 5kg and multi-layer bags containing 5kg and 25kg.



#### COMPOSITION

Each 100g contains:  
Ciprofloxacin HCl .....50g  
Excipient (enough. For) .....100g



#### ADMINISTRATION

Administer orally mixed in the feed or diluted in drinking water.



#### INDICATION

CIPROMAX it is indicated for the treatment of infections caused by Gram negative bacteria, such as Colibacillosis (*Escherichia coli*), Salmonellosis (*Salmonella typhimurium*; *S. enteritidis*) and by Gram positive bacteria, such as *Staphylococcus aureus* and *Streptococcus pyogenes*.



#### DOSAGE

Administer 10mg of ciprofloxacin per Kg of live weight for 3 5 consecutive days.



#### WAITING PERIOD

Suspend the use of the medication 2 days before the slaughter of animals intended for human consumption.





**MAXMOXIL**



#### PACKING

Sachets of 200g in boxes of 5kg and multi-layer bags containing 25kg.



#### COMPOSITION

Each 100g contains:  
Norfloxacin HCl.....50g  
Excipient (enough. For) .....100g



#### ADMINISTRATION

Administer orally mixed in the feed or diluted in drinking water.



#### INDICATION

MAXMOXIL is indicated for the treatment of bacterial infections caused by the bacteria: *Escherichia coli*, *Providencia retgeri* and *Citrobacter freundii*.



#### DOSAGE

Administer 7mg of norfloxacin per Kg of live weight for 5 7 consecutive days.



#### WAITING PERIOD

Suspend the use of the medication 5 days before the slaughter of animals intended for human consumption.



#### PACKING

Multilayer bags containing 25Kg.



#### INDICATION

OXIMAX it is indicated for the treatment of infections caused by *Escherichia coli*.



#### COMPOSITION

Each 100g contains:  
Oxytetracycline base.....50g  
Excipient (enough. For).....100g



#### DOSAGE

Administer 20mg to 30mg of oxytetracycline per Kg of body live weight for 3 to 5 consecutive days. It is recommended to continue treatment for 48 hours after the disappearance of symptoms.



#### ADMINISTRATION

For oral administration: mix with the feed.



#### WAITING PERIOD

Suspend the use of the medication 6 days before the slaughter of animals intended for human consumption.



## **DOXIVET 750**



### PACKING

Sachets of 100g in boxes of 5kg and multi-layer bags containing 5kg and 25kg.



### COMPOSITION

Each 100g contains:  
 Doxycycline (as hyclate) .....75g  
 Excipient (enough. For) .....100g



### ADMINISTRATION

Administer orally mixed in the feed or diluted in drinking water.



### INDICATION

DOXIVET 750 is indicated for the prevention and treatment of various diseases caused by agents sensitive to doxycycline, such as: *Bordetella bronchyseptica*, *Enterobacter aerogenes*, *Escherichia coli*, *Haemophilus sp.*, *Salmonella enteritidis*, and *Staphylococcus aureus*.



### DOSAGE

Administer 20mg of doxycycline per kg of body weight for 3 to 5 consecutive days. It is recommended to continue treatment for 48 hours after the disappearance of symptoms.



### WAITING PERIOD

Suspend the use of the medication 7 days before the sacrifice of the animals destined for human consumption if they were treated through the drinking water and if they were treated with the mixture in the feed: suspend 4 days before.



## DOXIVET 200



### PACKING

Sachets of 250g in boxes of 5kg and multilayer bags containing 25Kg.



### COMPOSITION

Each 100g contains:  
 Doxycycline HCl.....20g  
 Excipient (enough. For) .....100g



### ADMINISTRATION

It must be administered orally and mixed in the feed.



### INDICATION

DOXIVET 200 is indicated for the treatment and prevention of diseases caused by: *Haemophilus spp.*; *Mycoplasma spp.*; *Pasteurella spp.* and *Bordetella spp.* respiratory diseases caused by *Escherichia suis*, and *Actinobacillus pleuropneumoniae*; Enteric diseases caused by: *Escherichia coli*, *Clostridium spp.* and *Salmonella spp.*



### DOSAGE

Administer 500g to 1kg of DOXIVET 200 per ton of feed, this dose is equivalent to: 5 to 10mg/Kg of live weight, which is equivalent to: 100 to 200ppm of doxycycline per ton of feed. Continuous treatment from 5 to 10 days.



### WAITING PERIOD

Suspend the use of the medication 10 days before the slaughter of animals intended for human consumption.




**CURAMOXIL**



#### PACKING

Sachets of 200g in boxes of 5kg and multilayer bags containing 25Kg.



#### COMPOSITION

Each 100g contains:  
Amoxicillin Tri-Hydrate ..... 50g  
Excipient (enough. For) ..... 100g



#### ADMINISTRATION

Administer orally mixed in the feed or diluted in drinking water.



#### INDICATION

CURAMOXIL it is indicated for the treatment of infections caused by *coagulase negative Staphylococcus*, *Staphylococcus aureus*, *Klebsiella sp.*, *Pasteurella multocida* or *Pseudomonas aeruginos*, *Salmonella Typhimurim* and *Salmonella enteritidis*.



#### DOSAGE

Administer 20mg of amoxicillin per Kg live weight for 5 to 7 consecutive days. It is recommended to continue treatment for 48 hours after the disappearance of symptoms.



#### WAITING PERIOD

Suspend the use of the medication 1 day before the slaughter of the animals intended for human consumption.





## **AMOXMAX** **75% PREMIUM**



### PACKING

Multilayer bags containing 5kg and 25kg.



### INDICATION

AMOXMAX 75% PREMIUM is indicated for the treatment of infections caused by bacteria: *Escherichia coli*, *Salmonella*.



### COMPOSITION

Each 100g contains:  
 Amoxicillin (in Tri-hydrated form) ..... 75g  
 Excipient (enough. For) ..... 100g



### DOSAGE

Give 20mg of amoxicillin per kg of body live weight for 5 to 10 consecutive days. It is recommended to continue treatment for 48 hours after the disappearance of symptoms.



### ADMINISTRATION

Administer orally mixed with the feed.



### WAITING PERIOD

Suspend the use of the medication 2 days before the slaughter of animals intended for human consumption.



**FLORFENIMAX**



#### PACKING

Multilayer bags containing 25kg.



#### COMPOSITION

Each 100g contains:

Florfenicol.....4g

Excipient. (enough for) .....100g



#### ADMINISTRATION

Administer orally mixed with the feed.



#### INDICATION

FLORFENIMAX it is indicated for the treatment of respiratory tract diseases in pigs caused by: *Pasteurella multocida*, *Haemophilus parasuis* and *Escherichia coli*.



#### DOSAGE

Administer 2mg of florfenicol per kg of live weight for 7 to 14 consecutive days. It is recommended to continue treatment for 48 hours after the disappearance of symptoms.



#### WAITING PERIOD

Suspend the use of the medication 7 days before the slaughter of animals intended for human consumption.



## **FLORFENIMAX PS**



### PACKING

Sachets 200g in boxes of 5kg and multilayer bags containing 5kg and 25kg.



### INDICATION

FLORFENIMAX PS it is indicated for the treatment of pig infections caused by bacteria: *Pasteurella multocida*, *Haemophilus sp*, *Salmonella typhimurium*, *Escherichia coli* and *Erisipelothrix rhusiopathiae*.



### COMPOSITION

Each 100g contains:  
 Florfenicol..... 40g  
 Excipient. (enough for)..... 100g



### DOSAGE

Administer 15mg of florfenicol per Kg of live weight for 3 to 14 consecutive days.



### WAITING PERIOD

Suspend the use of the medication 6 days before the sacrifice of the animals destined for human consumption if they were treated through the drinking water and if they were treated with the mixture in the feed: suspend 4 days before.



### ADMINISTRATION

Administer orally mixed in the feed or diluted in drinking water.



**COLIMAX 50 PS**



#### PACKING

Sachets of 200g in boxes of 5kg and multilayer bags containing 5kg and 25kg.



#### INDICATION

COLIMAX 50 PS it is indicated for the treatment of enteric diseases caused by bacteria: *Escherichia coli* and *Pseudomonas aeruginosa*.



#### COMPOSITION

Each 100g contains:  
Colistin Sulfate .....50g  
Excipient (enough. For).....100g



#### DOSAGE

Administer 5mg of colistin sulfate per Kg of live weight for 5 consecutive days.



#### WAITING PERIOD

Suspend the use of the medication 2 days before the slaughter of animals intended for human consumption.



#### ADMINISTRATION

Administer orally mixed in the feed or diluted in drinking water.



## LINCOMAX 44



### PACKING

Sachets of 200g in boxes of 5kg and multi-layer bags containing 5kg and 25kg.



### COMPOSITION

Each 100g contains:  
Lincomycin (in the form of Lincomycin Hydrochloride) 44g  
Excipient (enough. For) ..... 100g



### ADMINISTRATION

Administer orally mixed in the feed or diluted in drinking water.



### INDICATION

LINCOMAX 44 is a medicine based on the antibiotic lincomycin, effective in combating lung and intestinal infections caused by bacteria: *Staphylococcus aureus* and *Streptococcus suis*.



### DOSAGE

Administer 5mg to 10mg of lincomycin per Kg of body weight for 7 to 10 consecutive days.



### WAITING PERIOD

Suspend the use of the medication 2 days before the slaughter of animals intended for human consumption.





  
**LINCOSPECMAX**



#### PACKING

Sachets of 100g in boxes of 5kg and multi-layer bags containing 5kg and 25kg.



#### COMPOSITION

Each 100g contains:

Lincomycin (in the form of Lincomycin Hydrochloride).....22g  
Spectinomycin (as Sulfate).....22g  
Excipient (enough. For) .....100g



#### INDICATION

LINCOSPECMAX is effective in combating lung and intestinal infections caused by bacteria: *Pasteurella multocida*, *Streptococcus suis*, *Salmonella enteritidis*, *Staphylococcus aureus*, *Salmonella typhimurium* and intestinal infections caused by: *Bordetella bronchiseptica*, *Klebsiella pneumoniae*, *Escherichia coli*, *Pasteurella multocida*, *Streptococcus suis*, *Pseudomonas aeruginosa*, *Proteus vulgaris*, *Salmonella enteritidis*, *Staphylococcus aureus* and *Salmonella typhimurium*.



#### DOSAGE

Administer 5mg to 10mg of the association per Kg of live weight for 5 to 10 consecutive days.



#### ADMINISTRATION

Administer orally mixed in the feed or diluted in drinking water.



#### WAITING PERIOD

Suspend the use of the medication 2 days before the slaughter of animals intended for human consumption.

**Stability in the Feed:** when mixed with the feed, it is stable for 30 days.

**Stability in Water:** When diluted in water, it is stable for 48 hours.

It is recommended to prepare a new mix after that period.

The presence of chlorine in the water does not interfere with the stability of the product.



#### PACKING

Sachets of 200g in boxes of 5kg and multi-layer bags containing 5kg and 25kg.



#### COMPOSITION

Each 100g contains:  
Neomycin (as Sulfate).....50g  
Excipient (enough. For) ..... 100g



#### ADMINISTRATION

Administer orally mixed in the feed or diluted in drinking water.



#### INDICATION

NEOMAX it is effective in combating infections caused by bacteria: *Clostridium perfringens*, *Escherichia coli*, *Klebsiella pneumoniae*, *Pasteurella multocida*, *Pseudomonas aeruginosa*, *Proteus mirabilis* and *Proteus vulgaris*, *Salmonella enteritidis* and *Salmonella typhimurium*.



#### DOSAGE

Administer 200g of neomycin for live weight for 7 consecutive days.



#### WAITING PERIOD

Suspend the use of the medication 2 days before the slaughter of animals intended for human consumption.



  
**CEFMAX**



#### PACKING

Glass bottle containing 100ml of the product.



#### COMPOSITION

Each 100g contains:  
Crystalline free acid ..... 10g  
Micronized ceftiofur ..... 10g  
Excipient. (enough for) ..... 100ml



#### ADMINISTRATION

Inject in the neck behind the ear, in a single dose. No more than 2 ml should be injected in a single application.  
Shake the bottle before use it.



#### INDICATION

CEFMAX it is indicated for the treatment and control of swine respiratory disease (DRS) and systemic disease, caused by: *Actinobacillus pleuropneumoniae*, *Streptococcus suis*, *Pasteurella multocida* and *Mannheimia haemolytica*.



#### DOSAGE

Administer a single dose of 5mg of ceftiofur per kg of body weight.



#### WAITING PERIOD

Suspend the use of the medication 2 days before the slaughter of animals intended for human consumption.



**RACTOMAX**



#### PACKING

Multilayer bags containing 5kg and 25kg.



#### COMPOSITION

Each 100g contains:  
 Ractopamine Hydrochloride .....2g  
 Excipient. (enough for) .....100g



#### ADMINISTRATION

Administer orally mixed with the feed.



#### INDICATION

RACTOMAX it is indicated for use exclusively in pigs in the final phase, to increase the rate of weight gain, improve feed efficiency and promote a lean carcass.



#### DOSAGE

Administer from 5ppm to 20ppm de ractopamine per ton of feed.



#### RETIREMENT PERIOD

This product does not require a withdrawal period for the slaughter of animals.



## ENRAPREMIUM PRO



### PACKING

Multilayer bags containing 25kg.



### COMPOSITION

Each 100g contains:

Enramycin ..... 8g  
Excipient (enough. For) ..... 100g



### ADMINISTRATION

Administer orally mixed with the feed.



### INDICATION

ENRAPREMIUM PRO it is a performance enhancing zootechnical additive based on Enramycin for manufacturers of pig feed products.



### DOSAGE

**Pigs (initial phase):** 62,5 to 125 g/ton of feed (equivalent to 5 to 10ppm of enramycin).  
**Pigs (growth/final phases):** 37,5 to 62,5g/ton of feed (equivalent to 3 to 5 ppm of enramycin).




### RETIREMENT PERIOD

This product does not require a withdrawal period for the slaughter of animals.





## FLUBENDAZOLE SAUVET



### PACKING

Multilayer bags containing 5kg and 25kg.



### COMPOSITION

Each 100g contains:  
 Flubendazole.....5g  
 Excipient. (enough for) .....100g



### ADMINISTRATION

Administer orally mixed with the feed.



### INDICATION

FLUBENDAZOLE SAUVET is effective for the treatment and control of gastrointestinal parasites: *Ascaris suum*, *Hyostrogylus rubidos*, *Oesophagostomum dentatum*, *Trichuris suis*, *Strongyloides ransomi* and pulmonary parasites: *Metastrongylus apri*.



### DOSAGE

Administer 5mg of flubendazole per Kg of live weight.  
**Sows and boars:** ½ sachet of 30g of FLUBENDAZOLE SAUVET for every 150kg of live weight.  
**Piglets:** ½ sachet of 30g of FLUBENDAZOLE SAUVET for a group of piglets with a weight equivalent to 150Kg.



### WAITING PERIOD

Suspend the use of the medication 15 days before the slaughter of animals intended for human consumption.





#### PACKING

1kg bottles containing 20g or 200g tablets.

#### COMPOSITION

Trichloro-s-triazine trione ..... 100%  
Active chlorine content ..... 90%

#### ADMINISTRATION

Administer diluted in drinking water.

#### INDICATION

MAXCLOR is a product for the treatment of drinking water, preventing the action of viruses, bacteria, fungi, and protozoa caused by *Staphylococcus aureus*, *Escherichia coli* and *Salmonella spp.*

#### DOSAGE

Administer the MAXCLOR tablets according to the water consumption demand of each shed or farm, to obtain the adequate free chlorine content (1 to 4ppm) in the drinker.

#### WAITING PERIOD

The product does not require a grace period before slaughtering the animals.

PRODUCT REGISTERED ON THE MAPA (Ministry of Agriculture and Livestock)



## **SANIMAX LINE**

### SANIMAX PLUS



#### COMPOSITION

Each 100ml contains:

Glutaraldehyde..... 42,5g  
Benzalkonium Chloride.....7,5g  
Excipient (enough. For).....100ml



#### ADMINISTRATION

**Disinfection of facilities and equipment:**

Dilute 1ml of SANIMAX PLUS in 1 to 2 liters of water.

**Disinfection of Incubators and slaughterhouses:**

Dilute 1ml of SANIMAX PLUS for each liter of water.

**Egg disinfection:**

Dilute 1ml of SANIMAX PLUS for each liter of water.

**Disinfection of footbaths, spray arches, vehicles:**

Dilute 1ml of SANIMAX PLUS for every 2 liters of water.



#### PACKING

Bottle containing 1 liter and bottle of 20 liters.



#### INDICATION

The SANIMAX line is indicated for the disinfection of incubators, slaughterhouses, eggs, egg grading machines, boxes for transporting and conditioning eggs, footbaths, vehicles and sheds.



### SANIMAX 800



#### COMPOSITION

Each 100ml contains:

Benzalkonium Chloride.....80g  
Excipient (enough. For).....100ml



#### ADMINISTRATION

Dilute in water in the proportion of 1 liter for every 1,000 liters of water or in accordance with the recommendation of the veterinarian.




### SANIMAX 500



#### COMPOSITION

Each 100ml contains:

Benzalkonium Chloride.....50g  
Excipient (enough. For).....100ml



Pharmacologically, the active ingredients used in the production of veterinary drugs can be presented in different forms, in their base or pure form, in the form of a salt or containing water molecules, and be in their hydrated form.

Considering these different forms, we must observe that the molecules of the active ingredients have an active fraction, which is responsible for the antimicrobial action, and an inactive fraction that does not exert antimicrobial action, even though they are often responsible for the stability, solubility, and palatability of the active ingredient.





The information on how the veterinary drug will be marketed will always be presented in the registration and label of the product. That is why SAUVET® always guides a careful verification of this information and in cases where there is a need for correction, observe if it is done at the time of drug production (legally allowed) or if it is necessary to use a correction factor, to reach the active fraction for an effective treatment.

To exemplify and show 100% of our quality and transparency, we attach to the table our entire portfolio and information on our products if they are corrected in their production (legally allowed) or if there is a need to use a factor of correction.


PRODUCTS  
**100%**  
QUALIFIED

In case of any doubt, our team is available to help our clients.



PRODUCT	ACTIVE INGREDIENTS	CONCENTRATION (%)	CORRECTION	SPECIES
 <b>Antimicrobials</b>				
<b>MAXIMULIN PS</b>	Tiamulin (Hydrogen fumarate)	45	Corrected	Poultry and Pigs
<b>MAXIMULIN 80</b>	Tiamulin (Hydrogen fumarate)	80	Corrected	Pigs
<b>MAXIMULIN 10% PREMIX</b>	Tiamulin (Hydrogen fumarate)	10	Corrected	Poultry and Pigs
<b>MEDMAXI</b>	Oxytetracycline Hydrochloride	10	Corrected	Poultry and Pigs
	Sulfadiazine base	7,5		
	Trimethoprim	1,5		
<b>ENROMAXI PS</b>	Enrofloxacin base	25	Corrected	Poultry
<b>MAXNOR 50</b>	Norfloxacin base	50	Corrected	Poultry and Pigs
<b>CIPROMAX</b>	Ciprofloxacin HCL	50	14,13%	Poultry and Pigs
<b>MAXMOXIL</b>	Norfloxacin HCL	50	10,26%	Poultry and Pigs
<b>OXIMAX</b>	Oxytetracycline base	50	Corrected	Poultry and Pigs
<b>DOXIVET 750</b>	Doxycycline (as hyclate)	75	Corrected	Poultry and Pigs
<b>DOXIVET 200</b>	Doxycycline HCl	20	13,36%	Poultry and Pigs
<b>CURAMOXIL</b>	Amoxicillin Tri-hydrate	50	12,88%	Poultry and Pigs
<b>AMOXMAX 75% PREMIUM</b>	Amoxicillin (in Tri-hydrated form)	75	Corrected	Poultry and Pigs
<b>FLORFENIMAX</b>	Florfenicol	4	Corrected	Pigs
<b>FLORFENIMAX PS</b>	Florfenicol	40	Corrected	Pigs
<b>LINCOMAX 44</b>	Lincomycin (as Lincomycin Hydrochloride)	44	Corrected	Poultry and Pigs
<b>LINCOSPECMAX</b>	Lincomycin (as Lincomycin Hydrochloride)	22	Corrected	Poultry and Pigs
	Spectinomycin (as Sulfate)	22		
<b>NEOMAX</b>	Neomycin (Sulfate)	50	Corrected	Poultry and Pigs
<b>CEFMAX</b>	Micronized Cefotiofur Crystalline Free Acid	10	Corrected	Pigs
 <b>Performance Enhancers</b>				
<b>RACTOMAX</b>	Ractopamine Hydrochloride	2	Corrected	Pigs
<b>ENRAPREMIUM PRO</b>	Enramycin	8	Corrected	Poultry and Pigs
<b>BACMAX</b>	Zinc Bacitracin	15	Corrected	Poultry
 <b>Antiparasitic</b>				
<b>FLUBENDAZOLE SAUVET</b>	Flubendazole	5	Corrected	Pigs
 <b>Biosecurity</b>				
<b>MAXCLOR</b>	Tricloro	90	Corrected	Poultry and Pigs
<b>SANIMAX 500</b>	Benzalkonium Chloride	50	Corrected	Poultry and Pigs
<b>SANIMAX 800</b>	Benzalkonium Chloride	80	Corrected	Poultry and Pigs
<b>SANIMAX PLUS</b>	Benzalkonium Chloride	7,5	Corrected	Poultry and Pigs
	Glutaraldehyde	42,5		



 Rua Antônio Lucchiari, 807 - Distrito Industrial  
Campinas / São Paulo - CEP 13054-700

 19 3846 3600

 [sauvet.com.br](http://sauvet.com.br)

 [facebook.com/sauvetbrasil](https://facebook.com/sauvetbrasil)

